arXiv:1007.3317v1 [math.NT] 20 Jul 2010

ON q-EULER NUMBERS RELATED TO THE MODIFIED q-BERNSTEIN POLYNOMIALS

MIN-SOO KIM, DAEYEOUL KIM AND TAEKYUN KIM

Abstract. We consider q-Euler numbers and polynomials and q-Stirling numbers of first and second kinds. Finally, we investigate some interesting properties of the modified q-Bernstein polynomials related to q-Euler numbers and q-Stirling numbers by using fermionic p-adic integrals on $\mathbb{Z}_p$.

1. Introduction

Let $C[0,1]$ be the set of continuous functions on $[0,1]$. The classical Bernstein polynomials of degree n for $f \in C[0,1]$ are defined by

$$\mathbb{B}_n(f) = \sum_{k=0}^{n} f\left(\frac{k}{n}\right) B_{k,n}(x), \quad 0 \le x \le 1 \tag{1.1}$$

where $\mathbb{B}_n(f)$ is called the Bernstein operator and

$$B_{k,n}(x) = \binom{n}{k} x^k (x-1)^{n-k} \tag{1.2}$$

are called the Bernstein basis polynomials (or the Bernstein polynomials of degree n) (see [17]). Recently, Acikgoz and Araci have studied the generating function for Bernstein polynomials (see [1, 2]). Their generating function for $B_{k,n}(x)$ is given by

$$F^{(k)}(t,x) = \frac{t^k e^{(1-x)t} x^k}{k!} = \sum_{n=0}^{\infty} B_{k,n}(x) \frac{t^n}{n!}, \tag{1.3}$$

where $k = 0, 1, \ldots$ and $x \in [0,1]$. Note that

$$B_{k,n}(x) = \begin{cases} \binom{n}{k} x^k (1-x)^{n-k} & \text{if } n \ge k \\ 0, & \text{if } n < k \end{cases}$$

for $n = 0, 1, \ldots$ (see [1, 2]).

Let p be an odd prime number. Throughout this paper, $\mathbb{Z}_p, \mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic rational integers, the field of p-adic rational numbers and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-1}$.

Throughout this paper, we use the following notation

$$[x]_q = \frac{1-q^x}{1-q} \quad \text{and} \quad [x]_{-q} = \frac{1-(-q)^x}{1+q}$$

2000 *Mathematics Subject Classification.* 11B68, 11S80.

Key words and phrases. q-Euler numbers and polynomials, q-Bernstein polynomials, q-Stirling numbers, Fermionic p-adic integrals.

(cf. [5, 6, 7, 12]). Let $\mathbb{N}$ be the natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable function on $\mathbb{Z}_p$.

Let $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-1/(p-1)}$ and $x \in \mathbb{Z}_p$. Then q-Bernstein type operator for $f \in UD(\mathbb{Z}_p)$ is defined by (see [14, 15])

$$\begin{aligned}\mathbb{B}_{n,q}(f) &= \sum_{k=0}^{n} f\left(\frac{k}{n}\right)\binom{n}{k}[x]_q^k[1-x]_q^{n-k} \\ &= \sum_{k=0}^{n} f\left(\frac{k}{n}\right) B_{k,n}(x,q),\end{aligned} \tag{1.4}$$

for $k, n \in \mathbb{Z}_+$, where

$$B_{k,n}(x,q) = \binom{n}{k}[x]_q^k[1-x]_q^{n-k} \tag{1.5}$$

is called the modified q-Bernstein polynomials of degree n. When we put $q \to 1$ in (1.5), $[x]_q^k \to x^k, [1-x]_q^{n-k} \to (1-x)^{n-k}$ and we obtain the classical Bernstein polynomial, defined by (1.2). We can deduce very easily from (1.5) that

$$B_{k,n}(x,q) = [1-x]_q B_{k,n-1}(x,q) + [x]_q B_{k-1,n-1}(x,q) \tag{1.6}$$

(see [14]). For $0 \leq k \leq n$, derivatives of the nth degree modified q-Bernstein polynomials are polynomials of degree $n-1$:

$$\frac{\mathrm{d}}{\mathrm{d}x} B_{k,n}(x,q) = n(q^x B_{k-1,n-1}(x,q) - q^{1-x} B_{k,n-1}(x,q))\frac{\ln q}{q-1} \tag{1.7}$$

(see [14]).

The Bernstein polynomials can also be defined in many different ways. Thus, recently, many applications of these polynomials have been looked for by many authors. In recent years, the q-Bernstein polynomials have been investigated and studied by many authors in many different ways (see [14, 15, 17] and references therein [4, 16]). In [16], Phillips gave many results concerning the q-integers, and an account of the properties of q-Bernstein polynomials. He gave many applications of these polynomials on approximation theory. In [1, 2], Acikgoz and Araci have introduced several type Bernstein polynomials. The Acikgoz and Araci paper to announce in the conference is actually motivated to write this paper. In [17], Simsek and Acikgoz constructed a new generating function of the q-Bernstein type polynomials and established elementary properties of this function. In [14], Kim, Jang and Yi proposed the modified q-Bernstein polynomials of degree n, which are different q-Bernstein polynomials of Phillips. In [15], Kim, Choi and Kim investigated some interesting properties of the modified q-Bernstein polynomials of degree n related to q-Stirling numbers and Carlitz's q-Bernoulli numbers.

In the present paper, we consider q-Euler numbers, polynomials and q-Stirling numbers of first and second kinds. We also investigate some interesting properties of the modified q-Bernstein polynomials of degree n related to q-Euler numbers and q-Stirling numbers by using fermionic p-adic integrals on $\mathbb{Z}_p$.

2. q-EULER NUMBERS AND POLYNOMIALS RELATED TO THE FERMIONIC p-ADIC INTEGRALS ON $\mathbb{Z}_p$

For $N \geq 1$, the fermionic q-extension μ_q of the p-adic Haar distribution μ_{Haar} :

$$\mu_{-q}(a + p^N\mathbb{Z}_p) = \frac{(-q)^a}{[p^N]_{-q}} \tag{2.1}$$

is known as a measure on $\mathbb{Z}_p$, where $a + p^N\mathbb{Z}_p = \{x \in \mathbb{Q}_p \mid |x - a|_p \leq p^{-N}\}$ (cf. [5, 8]). We shall write $d\mu_{-q}(x)$ to remind ourselves that x is the variable of integration. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable function on $\mathbb{Z}_p$. Then μ_{-q} yields the fermionic p-adic q-integral of a function $f \in UD(\mathbb{Z}_p)$:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N\to\infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} f(x)(-q)^x \tag{2.2}$$

(cf. [8, 9, 10, 13]). Many interesting properties of (2.2) were studied by many authors (see [8, 9] and the references given there). Using (2.2), we have the fermionic p-adic invariant integral on $\mathbb{Z}_p$ as follows:

$$\lim_{q\to-1} I_q(f) = I_{-1}(f) = \int_{\mathbb{Z}_p} f(a) d\mu_{-1}(x). \tag{2.3}$$

For $n \in \mathbb{N}$, write $f_n(x) = f(x + n)$. We have

$$I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2\sum_{l=0}^{n-1} (-1)^{n-l-1} f(l). \tag{2.4}$$

This identity is obtained by Kim in [8] to derives interesting properties and relationships involving q-Euler numbers and polynomials. For $n \in \mathbb{Z}_+$, we note that

$$I_{-1}([x]_q^n) = \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-1}(x) = E_{n,q}, \tag{2.5}$$

where $E_{n,q}$ are the q-Euler numbers (see [11]). It is easy to see that $E_{0,q} = 1$. For $n \in \mathbb{N}$, we have

$$\begin{aligned}
\sum_{l=0}^{n} \binom{n}{l} q^l E_{l,q} &= \sum_{l=0}^{n} \binom{n}{l} q^l \lim_{N\to\infty} \sum_{x=0}^{p^N-1} [x]_q^l (-1)^x \\
&= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} (-1)^x (q[x]_q + 1)^n \\
&= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} (-1)^x [x+1]_q^n \\
&= -\lim_{N\to\infty} \sum_{x=0}^{p^N-1} (-1)^x ([x]_q^n + [p^N]_q^n) \\
&= -E_{n,q}.
\end{aligned} \tag{2.6}$$

From this formula, we have the following recurrence formula

$$E_{0,q} = 1, \qquad (qE + 1)^n + E_{n,q} = 0 \quad \text{if } n \in \mathbb{N} \tag{2.7}$$

with the usual convention of replacing E^l by $E_{l,q}$. By the simple calculation of the fermionic p-adic invariant integral on $\mathbb{Z}_p$, we see that

$$E_{n,q} = \frac{2}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l \frac{1}{1+q^l}, \tag{2.8}$$

where $\binom{n}{l} = n!/l!(n-l)! = n(n-1)\cdots(n-l+1)/l!$. Now, by introducing the following equations:

$$[x]_{\frac{1}{q}}^n = q^n q^{-nx} [x]_q^n \quad \text{and} \quad q^{-nx} = \sum_{m=0}^{\infty} (1-q)^m \binom{n+m-1}{m} [x]_q^m \tag{2.9}$$

into (2.5), we find that

$$E_{n,\frac{1}{q}} = q^n \sum_{m=0}^{\infty} (1-q)^m \binom{n+m-1}{m} E_{n+m,q}. \tag{2.10}$$

This identity is a peculiarity of the p-adic q-Euler numbers, and the classical Euler numbers do not seem to have a similar relation. Let $F_q(t)$ be the generating function of the q-Euler numbers. Then we obtain

$$\begin{aligned} F_q(t) &= \sum_{n=0}^{\infty} E_{n,q} \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{2}{(1-q)^n} \sum_{l=0}^{n} (-1)^l \binom{n}{l} \frac{1}{1+q^l} \frac{t^n}{n!} \\ &= 2e^{\frac{t}{1-q}} \sum_{k=0}^{\infty} \frac{(-1)^k}{(1-q)^k} \frac{1}{1+q^k} \frac{t^k}{k!}. \end{aligned} \tag{2.11}$$

From (2.11) we note that

$$F_q(t) = 2e^{\frac{t}{1-q}} \sum_{n=0}^{\infty} (-1)^n e^{\left(\frac{-q^n}{1-q}\right)t} = 2\sum_{n=0}^{\infty} (-1)^n e^{[n]_q t}. \tag{2.12}$$

It is well-known that

$$I_{-1}([x+y]^n) = \int_{\mathbb{Z}_p} [x+y]^n d\mu_{-1}(y) = E_{n,q}(x), \tag{2.13}$$

where $E_{n,q}(x)$ are the q-Euler polynomials (see [11]). In the special case $x = 0$, the numbers $E_{n,q}(0) = E_{n,q}$ are referred to as the q-Euler numbers. Thus we have

$$\begin{aligned} \int_{\mathbb{Z}_p} [x+y]^n d\mu_{-1}(y) &= \sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} \int_{\mathbb{Z}_p} [y]^k d\mu_{-1}(y) \\ &= \sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} E_{k,q} \\ &= (q^x E + [x]_q)^n. \end{aligned} \tag{2.14}$$

It is easily verified, using (2.12) and (2.13), that the q-Euler polynomials $E_{n,q}(x)$ satisfy the following formula:

$$\begin{aligned}\sum_{n=0}^{\infty} E_{n,q}(x)\frac{t^n}{n!} &= \int_{\mathbb{Z}_p} e^{[x+y]_q t} d\mu_{-1}(y) \\ &= \sum_{n=0}^{\infty} \frac{2}{(1-q)^n} \sum_{l=0}^{n} (-1)^l \binom{n}{l} \frac{q^{lx}}{1+q^l} \frac{t^n}{n!} \qquad (2.15) \\ &= 2\sum_{n=0}^{\infty} (-1)^n e^{[n+x]_q t}.\end{aligned}$$

Using formula (2.15) when q tends to 1, we can readily derive the Euler polynomials, $E_n(x)$, namely,

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2e^{xt}}{e^t+1} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}$$

(see [8]). Note that $E_n(0) = E_n$ are referred to as the nth Euler numbers. Comparing the coefficients of $t^n/n!$ on both sides of (2.15), we have

$$E_{n,q}(x) = 2\sum_{m=0}^{\infty} (-1)^m [m+x]_q^n = \frac{2}{(1-q)^n} \sum_{l=0}^{n} (-1)^l \binom{n}{l} \frac{q^{lx}}{1+q^l}. \qquad (2.16)$$

We refer to $[n]_q$ as a q-integer and note that $[n]_q$ is a continuous function of q. In an obvious way we also define a q-factorial,

$$[n]_q! = \begin{cases} [n]_q[n-1]_q \cdots [1]_q & n \in \mathbb{N}, \\ 1, & n = 0 \end{cases}$$

and a q-analogue of binomial coefficient

$$\binom{x}{n}_q = \frac{[x]_q!}{[x-n]_q![n]_q!} = \frac{[x]_q[x-1]_q \cdots [x-n+1]_q}{[n]_q!} \qquad (2.17)$$

(cf. [10, 11]). Note that

$$\lim_{q\to 1} \binom{x}{n}_q = \binom{x}{n} = \frac{x(x-1)\cdots(x-n+1)}{n!}.$$

It readily follows from (2.17) that

$$\binom{x}{n}_q = \frac{(1-q)^n q^{-\binom{n}{2}}}{[n]_q!} \sum_{i=0}^{n} q^{\binom{i}{2}} \binom{n}{i}_q (-1)^{n+i} q^{(n-i)x} \qquad (2.18)$$

(cf. [11, 12]). It can be readily seen that

$$q^{lx} = ([x]_q(q-1)+1)^l = \sum_{m=0}^{l} \binom{l}{m} (q-1)^m [x]_q^m. \qquad (2.19)$$

Thus by (2.13) and (2.19), we have

$$\int_{\mathbb{Z}_p} \binom{x}{n}_q d\mu_{-1}(x) = \frac{(q-1)^n}{[n]_q! q^{\binom{n}{2}}} \sum_{i=0}^{n} q^{\binom{i}{2}} \binom{n}{i}_q (-1)^i \sum_{j=0}^{n-i} \binom{n-i}{j} (q-1)^j E_{j,q}. \qquad (2.20)$$

From now on, we use the following notation

$$\frac{[x]_q!}{[x-k]_q!}=q^{-\binom{k}{2}}\sum_{l=0}^{k}s_{1,q}(k,l)[x]_q^l,\quad k\in\mathbb{Z}_+,\tag{2.21}$$

$$[x]_q^n=\sum_{k=0}^{n}q^{\binom{k}{2}}s_{2,q}(n,k)\frac{[x]_q!}{[x-k]_q!},\quad n\in\mathbb{Z}_+\tag{2.22}$$

(see [12]). From (2.21), (2.22) and (2.19), we calculate the following consequence

$$\begin{aligned}
[x]_q^n&=\sum_{k=0}^{n}q^{\binom{k}{2}}s_{2,q}(n,k)\frac{1}{(1-q)^k}\sum_{l=0}^{k}\binom{k}{l}_q q^{\binom{l}{2}}(-1)^l q^{l(x-k+1)}\\
&=\sum_{k=0}^{n}q^{\binom{k}{2}}s_{2,q}(n,k)\frac{1}{(1-q)^k}\sum_{l=0}^{k}\binom{k}{l}_q q^{\binom{l}{2}+l(1-k)}(-1)^l\\
&\quad\times\sum_{m=0}^{l}\binom{l}{m}(q-1)^m[x]_q^m\\
&=\sum_{k=0}^{n}q^{\binom{k}{2}}s_{2,q}(n,k)\frac{1}{(1-q)^k}\\
&\quad\times\sum_{m=0}^{k}(q-1)^m\left(\sum_{l=m}^{k}\binom{k}{l}_q q^{\binom{l}{2}+l(1-k)}\binom{l}{m}(-1)^l\right)[x]_q^m.
\end{aligned}\tag{2.23}$$

Therefore, we obtain the following theorem.

Theorem 2.1. *For* $n\in\mathbb{Z}_+$,

$$E_{n,q}=\sum_{k=0}^{n}\sum_{m=0}^{k}\sum_{l=m}^{k}q^{\binom{k}{2}}s_{2,q}(n,k)(q-1)^{m-k}\binom{k}{l}_q q^{\binom{l}{2}+l(1-k)}\binom{l}{m}(-1)^{l+k}E_{m,q}.$$

By (2.19) and simple calculation, we find that

$$\begin{aligned}
\sum_{m=0}^{n}\binom{n}{m}(q-1)^m E_{m,q}&=\int_{\mathbb{Z}_p}q^{nx}d\mu_{-1}(x)\\
&=\sum_{k=0}^{n}(q-1)^k q^{\binom{k}{2}}\binom{n}{k}_q\int_{\mathbb{Z}_p}\prod_{i=0}^{k-1}[x-i]_q d\mu_{-1}(x)\\
&=\sum_{k=0}^{n}(q-1)^k\binom{n}{k}_q\sum_{m=0}^{k}s_{1,q}(k,m)\int_{\mathbb{Z}_p}[x]_q^m d\mu_{-1}(x)\\
&=\sum_{m=0}^{n}\left(\sum_{k=m}^{n}(q-1)^k\binom{n}{k}_q s_{1,q}(k,m)\right)E_{m,q}.
\end{aligned}\tag{2.24}$$

Therefore, we deduce the following theorem.

Theorem 2.2. *For* $n\in\mathbb{Z}_+$,

$$\sum_{m=0}^{n}\binom{n}{m}(q-1)^m E_{m,q}=\sum_{m=0}^{n}\sum_{k=m}^{n}(q-1)^k\binom{n}{k}_q s_{1,q}(k,m)E_{m,q}.$$

Corollary 2.3. *For $m, n \in \mathbb{Z}_+$ with $m \leq n$,*

$$\binom{n}{m}(q-1)^m = \sum_{k=m}^{n} (q-1)^k \binom{n}{k}_q s_{1,q}(k,m).$$

By (2.16) and Corollary 2.3, we obtain the following corollary.

Corollary 2.4. *For $n \in \mathbb{Z}_+$,*

$$E_{n,q}(x) = \frac{2}{(1-q)^n} \sum_{l=0}^{n} \sum_{k=l}^{n} (-1)^l (q-1)^{k-l} \binom{n}{k}_q s_{1,q}(k,l) \frac{q^{lx}}{1+q^l}.$$

It is easy to see that

$$\binom{n}{k}_q = \sum_{l_0+\cdots+l_k=n-k} q^{\sum_{i=0}^{k} i l_i} \tag{2.25}$$

(cf. [12]). From (2.25) and Corollary 2.4, we can also derive the following interesting formula for q-Euler polynomials.

Theorem 2.5. *For $n \in \mathbb{Z}_+$,*

$$E_{n,q}(x) = 2 \sum_{l=0}^{n} \sum_{k=l}^{n} \sum_{l_0+\cdots+l_k=n-k} q^{\sum_{i=0}^{k} i l_i} \frac{1}{(1-q)^{n+l-k}} s_{1,q}(k,l)(-1)^k \frac{q^{lx}}{1+q^l}.$$

These polynomials are related to the many branches of Mathematics, for example, combinatorics, number theory, discrete probability distributions for finding higher-order moments (cf. [10, 11, 13]). By substituting $x = 0$ into the above, we have

$$E_{n,q} = 2 \sum_{l=0}^{n} \sum_{k=l}^{n} \sum_{l_0+\cdots+l_k=n-k} q^{\sum_{i=0}^{k} i l_i} \frac{1}{(1-q)^{n+l-k}} s_{1,q}(k,l)(-1)^k \frac{1}{1+q^l}.$$

where $E_{n,q}$ is the q-Euler numbers.

3. q-EULER NUMBERS, q-STIRLING NUMBERS AND q-BERNSTEIN POLYNOMIALS RELATED TO THE FERMIONIC p-ADIC INTEGRALS ON $\mathbb{Z}_p$

First, we consider the q-extension of the generating function of Bernstein polynomials in (1.3).

For $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-1/(p-1)}$, we obtain

$$\begin{aligned} F_q^{(k)}(t,x) &= \frac{t^k e^{[1-x]_q t} [x]_q^k}{k!} \\ &= [x]_q^k \sum_{n=0}^{\infty} \binom{n+k}{k} [1-x]_q^n \frac{t^{n+k}}{(n+k)!} \\ &= \sum_{n=k}^{\infty} \binom{n}{k} [x]_q^k [1-x]_q^{n-k} \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} B_{k,n}(x,q) \frac{t^n}{n!}, \end{aligned} \tag{3.1}$$

which is the generating function of the modified q-Bernstein type polynomials (see [15]). Indeed, this generating function is also treated by Simsek and Acikgoz (see [17]). Note that $\lim_{q\to1} F_q^{(k)}(t,x)=F^{(k)}(t,x)$. It is easy to show that

$$[1-x]_q^{n-k}=\sum_{m=0}^{\infty}\sum_{l=0}^{n-k}\binom{l+m-1}{m}\binom{n-k}{l}(-1)^{l+m}q^l[x]_q^{l+m}(q-1)^m. \tag{3.2}$$

From (1.4), (2.3), (2.15) and (3.2), we derive the following theorem.

Theorem 3.1. *For $k,n\in\mathbb{Z}_+$ with $n\geq k$,*

$$\begin{aligned}\int_{\mathbb{Z}_p}&\frac{B_{k,n}(x,q)}{\binom{n}{k}}d\mu_{-1}(y)\\&=\sum_{m=0}^{\infty}\sum_{l=0}^{n-k}\binom{l+m-1}{m}\binom{n-k}{l}(-1)^{l+m}q^l(q-1)^mE_{l+m+k,q},\end{aligned}$$

where $E_{n,q}$ are the q-Euler numbers.

It is possible to write $[x]_q^k$ as a linear combination of the modified q-Bernstein polynomials by using the degree evaluation formulae and mathematical induction. Therefore we obtain the following theorem.

Theorem 3.2 ([14, Theorem 7]). *For $k,n\in\mathbb{Z}_+, i\in\mathbb{N}$ and $x\in[0,1]$,*

$$\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)=[x]_q^i([x]_q+[1-x]_q)^{n-i}.$$

Let $i-1\leq n$. Then from (1.5), (3.2) and Theorem 3.2, we have

$$\begin{aligned}[x]_q^i&=\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}\binom{n}{k}}{\binom{n}{i}}[x]_q^k[1-x]_q^{n-k}}{[x]_q^{n-i}\left(1+\frac{[1-x]_q}{[x]_q}\right)^{n-k}}\\&=\sum_{m=0}^{\infty}\sum_{k=i-1}^{n}\sum_{l=0}^{m+n-k}\sum_{p=0}^{\infty}\frac{\binom{k}{i}\binom{n}{k}}{\binom{n}{i}}\binom{l+p-1}{p}\binom{m+n-k}{l}\\&\quad\times\binom{n-i+m-1}{m}(-1)^{l+p+m}q^l(q-1)^p[x]_q^{i-n-m+k+p+l}.\end{aligned} \tag{3.3}$$

Using (2.13) and (3.3), we obtain the following theorem.

Theorem 3.3. *For $k,n\in\mathbb{Z}_+$ and $i\in\mathbb{N}$ with $i-1\leq n$,*

$$\begin{aligned}E_{i,q}=&\sum_{m=0}^{\infty}\sum_{k=i-1}^{n}\sum_{l=0}^{m+n-k}\sum_{p=0}^{\infty}\frac{\binom{k}{i}\binom{n}{k}}{\binom{n}{i}}\binom{l+p-1}{p}\binom{m+n-k}{l}\\&\times\binom{n-i+m-1}{m}(-1)^{l+p+m}q^l(q-1)^pE_{i-n-m+k+p+l,q}.\end{aligned}$$

The q-String numbers of the first kind is defined by

$$\prod_{k=1}^{n}(1+[k]_qz)=\sum_{k=0}^{n}S_1(n,k;q)z^k, \tag{3.4}$$

and the q-String number of the second kind is also defined by

$$\prod_{k=1}^{n}(1+[k]_q z)^{-1}=\sum_{k=0}^{n} S_2(n,k;q)z^k \tag{3.5}$$

(see [15]). Therefore, we deduce the following theorem.

Theorem 3.4 ([15, Theorem 4])**.** *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$,*

$$\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)}{([x]_q+[1-x]_q)^{n-i}}=\sum_{k=0}^{i}\sum_{l=0}^{k} S_1(n,l;q)S_2(i,k;q)[x]_q^l.$$

By Theorem 3.2, Theorem 3.4 and the definition of fermionic p-adic integrals on $\mathbb{Z}_p$, we obtain the following theorem.

Theorem 3.5. *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$,*

$$\begin{aligned} E_{i,q} &= \sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}\int_{\mathbb{Z}_p}\frac{B_{k,n}(x,q)}{([x]_q+[1-x]_q)^{n-i}}d\mu_{-1}(x) \\ &= \sum_{k=0}^{i}\sum_{l=0}^{k} S_1(n,l;q)S_2(i,k:q)E_{l,q}, \end{aligned}$$

where $E_{i,q}$ is the q-Euler numbers.

Let $i-1 \le n$. It is easy to show that

$$\begin{aligned} & [x]_q^i([x]_q+[1-x]_q)^{n-i} \\ &= \sum_{l=0}^{n-i}\binom{n-i}{l}[x]_q^{l+i}[1-x]_q^{n-i-l} \\ &= \sum_{l=0}^{n-i}\sum_{m=0}^{n-i-l}\binom{n-i}{l}\binom{n-i-l}{m}(-1)^m q^m [x]_q^{m+i+l} q^{-mx} \\ &= \sum_{l=0}^{n-i}\sum_{m=0}^{n-i-l}\sum_{s=0}^{\infty}\binom{n-i}{l}\binom{n-i-l}{m}\binom{m+s-1}{s} \\ &\quad \times (-1)^m q^m (1-q)^s [x]_q^{m+i+l+s}. \end{aligned} \tag{3.6}$$

From (3.6) and Theorem 3.2, we have the following theorem.

Theorem 3.6. *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$,*

$$\begin{aligned} \sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}\int_{\mathbb{Z}_p}B_{k,n}(x,q)d\mu_{-1}(x) &= \sum_{l=0}^{n-i}\sum_{m=0}^{n-i-l}\sum_{s=0}^{\infty}\binom{n-i}{l}\binom{n-i-l}{m}\binom{m+s-1}{s} \\ &\quad \times (-1)^m q^m (1-q)^s E_{m+i+l+s,q}, \end{aligned}$$

where $E_{i,q}$ are the q-Euler numbers.

In the same manner, we can obtain the following theorem.

Theorem 3.7. *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$,*

$$\begin{aligned}\int_{\mathbb{Z}_p} B_{k,n}(x,q)\,d\mu_{-1}(x) = &\sum_{j=k}^{n}\sum_{m=0}^{\infty}\binom{j}{k}\binom{n}{j}\binom{j-k+m-1}{m}\\ &\times(-1)^{j-k+m}q^{j-k}(q-1)^m E_{m+j,q},\end{aligned}$$

where $E_{i,q}$ are the q-Euler numbers.

4. Further remarks and observations

The q-binomial formulas are known,

$$\begin{aligned}(a;q)_n &= (1-a)(1-aq)\cdots(1-aq^{n-1}) = \sum_{i=0}^{n}\binom{n}{i}_q q^{\binom{i}{2}}(-1)^i a^i,\\ \frac{1}{(a;q)_n} &= \frac{1}{(1-a)(1-aq)\cdots(1-aq^{n-1})} = \sum_{i=0}^{\infty}\binom{n+i-1}{i}_q a^i.\end{aligned}\tag{4.1}$$

For $h \in \mathbb{Z}, n \in \mathbb{Z}_+$ and $r \in \mathbb{N}$, we introduce the extended higher-order q-Euler polynomials as follows [11]:

$$E_{n,q}^{(h,r)}(x) = \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p} q^{\sum_{j=1}^{r}(h-j)x_j}[x+x_1+\cdots+x_r]_q^n\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r).\tag{4.2}$$

Then

$$\begin{aligned}E_{n,q}^{(h,r)}(x) &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l\frac{q^{lx}}{(-q^{h-1+l};q^{-1})_r}\\ &= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l\frac{q^{lx}}{(-q^{h-r+l};q)_r}.\end{aligned}\tag{4.3}$$

Let us now define the extended higher-order Nörlund type q-Euler polynomials as follows [11]:

$$\begin{aligned}E_{n,q}^{(h,-r)}(x) = &\frac{1}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l\\ &\times\frac{q^{lx}}{\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p} q^{l(x_1+\cdots+x_r)}q^{\sum_{j=1}^{r}(h-j)x_j}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)}.\end{aligned}\tag{4.4}$$

In the special case $x=0$, $E_{n,q}^{(h,-r)} = E_{n,q}^{(h,-r)}(0)$ are called the extended higher-order Nörlund type q-Euler numbers. From (4.4), we note that

$$\begin{aligned}E_{n,q}^{(h,-r)}(x) &= \frac{1}{2^r(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{lx}(-q^{h-r+l};q)_r\\ &= \frac{1}{2^r}\sum_{m=0}^{r} q^{\binom{m}{2}}q^{(h-r)m}\binom{r}{m}_q[m+x]_q^n.\end{aligned}\tag{4.5}$$

A simple manipulation shows that

$$q^{\binom{m}{2}}\binom{r}{m}_q = \frac{q^{\binom{m}{2}}[r]_q\cdots[r-m+1]_q}{[m]_q!} = \frac{1}{[m]_q!}\prod_{k=0}^{m-1}([r]_q-[k]_q)\tag{4.6}$$

and

$$\prod_{k=0}^{n-1}(z-[k]_q)=z^n\prod_{k=0}^{n-1}\left(1-\frac{[k]_q}{z}\right)=\sum_{k=0}^{n}S_1(n-1,k;q)(-1)^k z^{n-k}. \tag{4.7}$$

Formulas (4.5), (4.6) and (4.7) imply the following lemma.

Lemma 4.1. *For $h\in\mathbb{Z}, n\in\mathbb{Z}_+$ and $r\in\mathbb{N}$,*

$$E_{n,q}^{(h,-r)}(x)=\frac{1}{2^r[m]_q!}\sum_{m=0}^{r}\sum_{k=0}^{m}q^{(h-r)m}S_1(m-1,k;q)(-1)^k[r]_q^{m-k}[x+m]_q^n.$$

From (2.19), we can easily see that

$$[x+m]_q^n=\frac{1}{(1-q)^n}\sum_{j=0}^{n}\sum_{l=0}^{j}\binom{n}{j}\binom{j}{l}(-1)^{j+l}(1-q)^l q^{mj}[x]_q^l. \tag{4.8}$$

Using (2.13) and (4.8), we obtain the following lemma.

Lemma 4.2. *For $m,n\in\mathbb{Z}_+$,*

$$E_{n,q}(m)=\frac{1}{(1-q)^n}\sum_{j=0}^{n}\sum_{l=0}^{j}\binom{n}{j}\binom{j}{l}(-1)^{j+l}(1-q)^l q^{mj}E_{l,q}.$$

By Lemma 4.8, lemma 4.2 and the definition of fermionic p-adic integrals on $\mathbb{Z}_p$, we obtain the following theorem.

Theorem 4.3. *For $h\in\mathbb{Z}, n\in\mathbb{Z}_+$ and $r\in\mathbb{N}$,*

$$\begin{aligned}\int_{\mathbb{Z}_p}E_{n,q}^{(h,-r)}(x)d\mu_{-1}(x)&=\frac{2^{-r}}{[m]_q!}\sum_{m=0}^{r}\sum_{k=0}^{m}q^{(h-r)m}S_1(m-1,k;q)(-1)^k[r]_q^{m-k}E_{n,q}(m)\\&=\frac{1}{2^r[m]_q!}\sum_{m=0}^{r}\sum_{k=0}^{m}q^{(h-r)m}S_1(m-1,k;q)(-1)^k[r]_q^{m-k}\\&\quad\times\frac{1}{(1-q)^n}\sum_{j=0}^{n}\sum_{l=0}^{j}\binom{n}{j}\binom{j}{l}(-1)^{j+l}(1-q)^l q^{mj}E_{l,q}.\end{aligned}$$

Put $h=0$ in (4.4). We consider the following polynomials $E_{n,q}^{(0,-r)}(x)$:

$$E_{n,q}^{(0,-r)}(x)=\sum_{l=0}^{n}\frac{(1-q)^{-n}\binom{n}{l}(-1)^l q^{lx}}{\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{l(x_1+\cdots+x_r)}q^{-\sum_{j=1}^{r}jx_j}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)}. \tag{4.9}$$

Then

$$E_{n,q}^{(0,-r)}(x)=\frac{1}{2^r}\sum_{m=0}^{r}\binom{r}{m}q^{\binom{m}{2}-rm}[m+x]_q^n.$$

A simple calculation of the ferminionic p-adic invariant integral on $\mathbb{Z}_p$ shows that

$$\int_{\mathbb{Z}_p}E_{n,q}^{(0,-r)}(x)d\mu_{-1}(x)=\frac{1}{2^r}\sum_{m=0}^{r}\binom{r}{m}q^{\binom{m}{2}-rm}E_{n,q}(m).$$

Using Theorem 4.3, we can also prove that

$$\int_{\mathbb{Z}_p}E_{n,q}^{(0,-r)}(x)d\mu_{-1}(x)=\frac{2^{-r}}{[m]_q!}\sum_{m=0}^{r}\sum_{k=0}^{m}q^{-rm}S_1(m-1,k;q)(-1)^k[r]_q^{m-k}E_{n,q}(m).$$

Therefore, we obtain the following theorem.

Theorem 4.4. *For $m \in \mathbb{Z}_+, r \in \mathbb{N}$ with $m \leq r$,*

$$\binom{r}{m} q^{\binom{m}{2}-rm} = \frac{1}{[m]_q!} \sum_{k=0}^{m} q^{-rm} S_1(m-1,k;q)(-1)^k [r]_q^{m-k}.$$

References

- [1] M. Acikgoz and S. Araci, *A study on the integral of the product of several type Bernstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, 2010.
- [2] M. Acikgoz and S. Araci, *On the generating function of the Bernstein polynomials*, Accepted to AIP on 24 March 2010 for ICNAAM 2010.
- [3] S. Bernstein, *Demonstration du theoreme de Weierstrass, fondee sur le calcul des probabilities*, Commun. Soc. Math. Kharkow (2) **13** (1912-1913), 1–2.
- [4] N. K. Govil and V. Gupta, *Convergence of q-Meyer-Konig-Zeller-Purrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 181–189.
- [5] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320–329.
- [6] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), no. 3, 288–299.
- [7] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12** (2005), 186–196.
- [8] T. Kim, *On the analogs of Euler numbers and polynomials associated with p-adic q-integral on $\mathbb{Z}_p$ at $q = -1$*, J. Math. Anal. Appl. **331** (2007), 779–792.
- [9] T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), 51–57.
- [10] T. Kim, *On p-adic interpolating function for q-Euler numbers and its derivatives*, J. Math. Anal. Appl. **339** (2008), 598–608.
- [11] T. Kim, *Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. **16** (2009), no. 4, 484–491.
- [12] T. Kim, *q-Bernoulli numbers associated with q-Stirling numbers*, Adv. Difference Equ. 2008, Art. ID 743295, 10 pp.
- [13] T. Kim, *Barnes-type multiple q-zeta functions and q-Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201.
- [14] T. Kim, L.-C. Jang and H. Yi, *Note on the modified q-Bernstein polynomials*, Discrete Dynamics in Nature and Society (in Press). arXiv 1005.4293.
- [15] T. Kim, J. Choi and Y.-H. Kim *Some identities on the q-Bernstein polynomials, q-Stirling numbers and q-Bernoulli numbers*, Adv. Stud. Contemp. Math. 20 (2010), no. 3, 335–341. arXiv 1006.2033.
- [16] G. M. Phillips, *Bernstein polynomials based on the q-integer*, Annals of Numerical Analysis **4** (1997), 511–518.
- [17] Y. Simsek and M. Acikgoz, *A new generating function of q-Bernstein-type polynomials and their interpolation function*, Abstr. Appl. Anal. ID 769095 (2010), 12 pp.

Department of Mathematics, KAIST, 373-1 Guseong-dong, Yuseong-gu, Daejeon 305-701, S. Korea
E-mail address: `minsookim@kaist.ac.kr`

National Institute for Mathematical Sciences, Doryong-dong, Yuseong-gu, Daejeon 305-340, Republic of KOREA
E-mail address: `daeyeoul@nims.re.kr`

Division of General Education-Mathematics, Kwangwoon University, Seoul, 139-701, S. Korea
E-mail address: `tkkim@kw.ac.kr`